

R. Kainhofer, R. Tichy

Graz University of Technology

**Quasi-randomized schemes for the
solution of retarded differential
equations**

Talk given at the Dagstuhl Seminar 2401
"Algorithms and Complexity for Continuous
Problems"

Schloss Dagstuhl, September 29 - October
4, 2002

Overview

1. Sketch of the numerical solution
2. Historic background
3. The RKQMC solution methods (Hermite Interpolation, QMC methods)
4. Convergence proofs
5. Numerical examples

The problem

Heavily varying delay differential equations (DDE) or DDE with heavily varying solutions.

$$y'(t) = f(t, y(t), y(t - \tau_1(t)), \dots, y(t - \tau_k(t))),$$

for $t \geq t_0$, $k \geq 1$,

$$y(t) = \varphi(t), \text{ for } t \leq t_0,$$

with

$f(t, \cdot, \cdot)$...piecewise smooth in y and y_{ret} ,
only bounded and Borel measurable in t
 $y(t)$...solution, d -dimensional real-valued function
 $\tau_1(t), \dots, \tau_k(t)$...continuous delay functions,
bounded from below by $\tau_0 > 0$,
satisfy $t_1 - \tau_j(t_1) \leq t_2 - \tau_j(t_2)$ for $t_1 \leq t_2$
 $\varphi(t)$...initial function, continuous at least
on $\left[\inf_{\substack{t_0 \leq t \\ 1 \leq j \leq k}} (t - \tau_j(t)), t_0 \right]$.

For heavily oscillating DDE: conventional Runge-Kutta methods unstable.

Sketch of the numerical solution

- Hermite interpolation for retarded argument \Rightarrow ODE
- use RKQMC methods for ODE:
Large Runge-Kutta error for heavily oscillating DE \rightarrow Idea (Stengle, Lécot):
integrate over whole step size
 - Runge Kutta: Integration over y and t discretized
 - RK(Q)MC: Integration over y discretized, numerical Integration in t (using MC or QMC integration to minimize the error)

Historical background

Halton (1960), Sobol (1960/67), Niederreiter (1978), Faure (1981/82):

Low-discrepancy sequences

Oppelstrup (1976), Oberle and Pesch (1981), ...:

Treatment of DDE with Hermite interpolation for retarded arguments

Stengle (1990, 1995):

Randomized Runge Kutta schemes for (heavily oscillating) ordinary differential equations

Coulibaly and Lécot (1999), Lécot (2001), Lécot and Koudiraty (2002):

Quasi-Monte Carlo Runge Kutta schemes (using low-discrepancy sequences) for ODE

Kainhofer and Tichy (2001/02):

RKQMC methods for delayed differential equations, combining the Hermite interpolation with RKQMC methods.

Hermite interpolation

Use hermite interpolation for the retarded arguments:

$$z_j(t) := y(t - \tau_j(t)) = \begin{cases} \varphi(t - \tau_j(t)), & \text{if } t - \tau_j(t) \leq t_0 \\ P_q(t - \tau_j(t); (y_i); (y'_i)) & \text{otherwise} \end{cases}$$

DDE transforms to a ODE:

$$\begin{aligned} y'(t) &= f(t, y(t), y(t - \tau_1(t)), \dots, y(t - \tau_k(t))) \approx \\ &\approx f(t, y(t), z_1(t), \dots, z_k(t)) =: g(t, y(t)) . \end{aligned}$$

Solution has to be piecewise $r/2$ -times continuously differentiable in t .

Resulting ODE has to fulfill requirements for RKQMC methods (Borel-measurable in t , continuously differentiable in $y(t)$).

Runge Kutta Quasi-Monte Carlo methods

G. Stengle, Ch. Lécot, I. Coulibaly, A. Koudiry

$$\begin{aligned}y'(t) &= f(t, y(t)), & 0 < t < T, \\y(0) &= y_0\end{aligned}$$

f smooth in y , bounded and Borel measurable in t .

f is Taylor-expanded only in y
→ integral equation in t .

e.g. 2nd order:

$$y(t_{n+1}) = y(t_n) + \frac{1}{2h_n} \int_{t_n}^{t_{n+1}} \int_{t_n}^{t_{n+1}} \left(f(\underline{u}, y(t_n)) + \frac{1}{\beta} f(\bar{u}, y(t_n)) + \frac{1}{\alpha} f(\bar{u}, y(t_n) + \alpha h_n f(\underline{u}, y(t_n))) \right) du$$

Solved by (Quasi-)Monte Carlo integration

Quasi-Monte Carlo methods

Integrals replaced by a discrete sum over N (quasi-)random points:

$$\int_{[0,1]^s} f(x) dx = \frac{1}{N} \sum_{i=1}^N f(x_i)$$

MC methods: x_i random points

QMC methods: x_i low-discrepancy sequences

Low discrepancy sequences: deterministic point sequences $\{x_i\}_{1 \leq i \leq N} \in [0, 1]^s$, good uniform distribution

star-discrepancy of the point set S

$$D_N^*(S) = \sup_{a,b \in [0,1]^s} \left| \frac{A([0, 1]^s, S)}{N} - \lambda_s([a, b]) \right|$$


Koksma-Hlawka inequality:

$$\left| \frac{1}{N} \sum_{n=1}^N f(x_n) - \int_{[0,1]^s} f(u) du \right| \leq V([0, 1]^s, f) D_N^*(x_1, \dots, x_N) .$$

Low-Discrepancy sequences

$$D_N^*(S) \leq \mathcal{O}\left(\frac{(\log N)^s}{N}\right)$$

- Halton-sequence in bases (b_1, \dots, b_s) : inversion of digit expansion of n in base b_i at the comma
- (t, s) nets in base b (Niederreiter, Sobol, Faure): net-like structure \rightarrow best possible uniform distribution on elementary intervals



Problem: correlations between elements

general form of RKQMC method

$$y_{n+1} = y_n + \frac{h_n}{s!N} \sum_{0 \leq j < N} G_s(\bar{t}_{j,n}; y)$$

$G_s(\bar{t}_{j,n}; y) \dots$ differential increment function of scheme

$$G_1(u; y) = f(u, y)$$

$$G_2(\bar{u}; y) = f(\bar{u}_1, y) + \frac{1}{\beta} f(\bar{u}_2, y) + \frac{1}{\alpha} f(\bar{u}_2, y + \alpha h_n f(\bar{u}_1, y))$$

$$G_3(\bar{u}; y) = a_1 f(\bar{u}_1, y) + \sum_{l=1}^{L_2} a_{2,l} f(\bar{u}_2, y + b_{2,l} h_n f(\bar{u}_1, y)) + \\ + \sum_{l=1}^{L_3} a_{3,l} (\bar{u}_3, y + b_{3,l}^{(1)} h_n f(\bar{u}_1, y) + b_{3,l}^{(2)} h_n f(\bar{u}_2, y + c_{3,l} h_n f(\bar{u}_1, y)))$$

Convergence proof by Koudiraty and Lécot:

If initial error $\|e_0\|$, the step size H , the discrepancy $D_N^*(X)$ and $\|f\|_E$ are small enough,

$$\|e_n\| \leq e^{c_2 t_n} \|e_0\| + \frac{e^{c_2 t_n} - 1}{c_2} (c_1 H^3 + c_3 D_N^*(X))$$

RKQMC for Volterra functional equations

$$\begin{aligned}y'(t) &= f(t, y(t), z(t)) & t \geq t_0 \\z(t) &= (Fy)(t) \\y(s) &= \varphi(s) & s \leq t_0\end{aligned}$$

- $(Fy)(t)$ contains dependence on retarded values.
- F and f are Lipschitz continuous in all arguments except t .

RKQMC method ($n \geq 0$):

$$\begin{aligned}y_{n+1} &= y_n + h_n \sum_{i=1}^N G_s \left(t_{n,i}; y_n, \tilde{z}(t) \right) \\ \tilde{z}(t) &= (\tilde{F}y_j)(t)\end{aligned}$$

Convergence theorems: RKQMC for DDE

Theorem: (Kainhofer, 2002)

If

- (i) RKQMC method converges for ordinary differential equations with order p
- (ii) the increment function G_s of the method and F are Lipschitz
- (iii) the interpolation fulfills a Lipschitz condition
- (iv) Hermite interpolation is used with order r
- (v) the initial error $\|e_0\|$ vanishes

then the method converges. If at least ii and iii hold, the error is bounded by

$$\|e_{j+1}\| \leq \|e_0\| e^{\mathcal{L}t_j} + \frac{(e^{t_j\mathcal{L}} - 1)}{\mathcal{L}} (\|E_G^{\text{ODE}}\| + \mathcal{L}_1\mathcal{L}_2 \|E_r^{\text{interpol}}\|) .$$

Special case: one retarded argument

Choose the RKQMC method:

- G_s Lipschitz (\mathcal{L}_2) in 2nd and 3rd argument, bounded variation (in the sense of Hardy and Krause).

- $\exists c_1, c_2, c_3$ such that for a $p > 0$

$$\text{loc. trunc. error } \|\epsilon_n\| \leq c_1(h_n)h_n^p$$

$$\text{RK error } \|\delta_n\| \leq c_2(h_n) \|e_n\|$$

$$\text{QMC error } \|d_n\| \leq c_3(h_n)D_N^*(S)$$

- interpolation order p , fulfils a certain Lipschitz condition

Then the error $\|e_n\| = \|y_n - y(t_n)\|$ of the method is bounded from above by:

$$\|e_n\| \leq \|e_0\| e^{t_n \left(c_2 + \frac{\mathcal{L}_2}{s!} \right)} + \frac{e^{t_n \left(c_2 + \frac{\mathcal{L}_2}{s!} \right)} - 1}{c_2 + \frac{\mathcal{L}_2}{s!}} \cdot \left\{ c_3 D_N^*(X) + \frac{\mathcal{L}_2}{s!} M H^q + c_1 H^p \right\}$$

Numerical examples

$$y'(t) = 3y(t-1)\sin(\lambda t) + 2y(t-1.5)\cos(\lambda t), \quad t \geq 0$$

$$y(t) = 1, \quad t \leq 0,$$

$h_n = 0.001$ for RK, $h_n = 0.01$ for RKQMC

i




Fig: Error of the RK and RKQMC methods

slowly varying (small λ):
conventional RK better

rapidly varying (high λ):

RKQMC outperform higher order RK

Advantage of RKQMC for unstable DDE

$$y'(t) = \pi \frac{\lambda}{2} \left(y \left(t - 2 - \frac{3}{2\lambda} \right) - y \left(t - 2 - \frac{1}{2\lambda} \right) \right), \quad t \geq 0$$

$$y(t) = \sin(\lambda t \pi), \quad t < 0,$$

Exact Solution: $y(t) = \sin(\lambda t \pi)$

- No discontinuities in any derivative
- k -th derivative only bounded by $\lambda^k \rightarrow$ "exploding" error

Numerical results:




Fig: Error of the RK and RKQMC methods

RKQMC schemes can delay the instability of the solution for heavily oscillating delay differential equations.

Time-corrected error

QMC integration is more expensive than 1 evaluation (Runge-Kutta)

→ Compare a time-corrected error

Result: RKQMC methods loose some advantage, but still better than Runge-Kutta for heavily oscillating DDE.




Fig: Time-corrected error of the RK and RKQMC methods

References

1. I. Coulibaly and C. Lécot. *A quasi-randomized Runge-Kutta method*. *Mathematics of Computation*, 68(226):651–659, 1999.
2. R. Kainhofer. *QMC Methods for the solution of delay differential equations*. Preprint, submitted 2002.
3. R. Kainhofer and R.F. Tichy. *QMC Methods for the Solution of Differential Equations with Multiple Delayed Arguments*. *Grazer Math. Berichte*, 2002 (to appear).
4. A. A. Koudiraty. *Numerical analysis of some quasi-Monte Carlo Methods*. PhD thesis, Université de Savoie, 2001.
5. C. Lécot. *Quasi-randomized numerical methods for systems with coefficients of bounded variation*. *Mathematics and Computers in Simulation*, 55:113–121, 2001.
6. H. J. Oberle and H. J. Pesch. *Numerical Treatment of Delay Differential Equations by Hermite Interpolation*. *Numer. Math.*, 37:235–255, 1981.
7. G. Stengle. *Error analysis of a randomized numerical method*. *Numer. Math.*, 70:119–128, 1995.